


**NEET(UG)-2018 TEST PAPER WITH ANSWER & SOLUTION  
 (HELD ON SUNDAY 06<sup>th</sup> MAY, 2018)**
**CHEMISTRY**

46. A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc.  $H_2SO_4$ . The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be  
 (1) 1.4 (2) 3.0 (3) 2.8 (4) 4.4


**Ans. (3)**

**Sol.**



$$(\text{moles})_i = \frac{2.3}{46} = \frac{1}{20} \quad 0 \quad 0$$

$$(\text{moles})_f \quad 0 \quad \frac{1}{20} \quad \frac{1}{20}$$



$$(\text{moles})_i \quad \frac{4.5}{90} = \frac{1}{20} \quad 0 \quad 0 \quad 0$$

$$(\text{moles})_f \quad 0 \quad \frac{1}{20} \quad \frac{1}{20} \quad \frac{1}{20}$$

$CO_2$  is absorbed by KOH.

So the remaining product is only CO.

moles of CO formed from both reactions

$$= \frac{1}{20} + \frac{1}{20} = \frac{1}{10}$$

Left mass of CO = moles  $\times$  molar mass


$$= \frac{1}{10} \times 28$$

$$= \boxed{2.8 \text{ g}} \text{ Ans.}$$

47. Nitration of aniline in strong acidic medium also gives m-nitroaniline because  
 (1) In spite of substituents nitro group always goes to only m-position.  
 (2) In electrophilic substitution reactions amino group is meta directive.  
 (3) In absence of substituents nitro group always goes to m-position  
 (4) In acidic (strong) medium aniline is present as anilinium ion.

**Ans. (4)**

**Sol.**



In acidic medium aniline is protonated to form anilinium ion which is metadirecting.

48. Which of the following oxides is most acidic in nature?

- (1) MgO (2) BeO  
 (3) BaO (4) CaO

**Ans. (2)**

**Sol.** In metals moving down the group metallic character increases, so basic nature increases hence most acidic will be BeO.

49. The difference between amylose and amylopectin is

- (1) Amylopectin have 1  $\rightarrow$  4  $\alpha$ -linkage and 1  $\rightarrow$  6  $\alpha$ -linkage  
 (2) Amylose have 1  $\rightarrow$  4  $\alpha$ -linkage and 1  $\rightarrow$  6  $\beta$ -linkage  
 (3) Amylopectin have 1  $\rightarrow$  4  $\alpha$ -linkage and 1  $\rightarrow$  6  $\beta$ -linkage  
 (4) Amylose is made up of glucose and galactose

**Ans. (1)**

**Sol.**

Amylose is long unbranched chain with  $\alpha$ -D-Glucose with held by  $C_1$ - $C_4$  glucosidic linkage whereas amylopectin is branched chain polymer of  $\alpha$ -D glucose unit in which chain is formed by  $C_1$ - $C_4$  glycosidic linkage while branching occurs by  $C_1$ - $C_6$  glucosidic linkage.


50. Regarding cross-linked or network polymers, which of the following statements is **incorrect**?

- (1) They contain covalent bonds between various linear polymer chains.  
 (2) They are formed from bi-and tri-functional monomers.  
 (3) Examples are bakelite and melamine.  
 (4) They contain strong covalent bonds in their polymer chains.

**Ans. (4)**

**Sol.** Cross-linked or network polymers are usually formed from bi-functional & tri-functional monomers and contains strong covalent bond between various linear polymer chains like Melamine, Bakelite etc.

51. In the reaction




the electrophile involved is

- (1) dichloromethyl cation ( $\text{CHCl}_2^+$ )
- (2) formyl cation ( $\text{CHO}^+$ )
- (3) dichloromethyl anion ( $\text{CHCl}_2^-$ )
- (4) dichlorocarbene ( $:\text{CCl}_2$ )

Ans. (4)

Sol.




52. Carboxylic acid have higher boiling points than aldehydes, ketones and even alcohols of comparable molecular mass. It is due to their

- (1) formation of intramolecular H-bonding
- (2) formation of carboxylate ion
- (3) more extensive association of carboxylic acid via van der Waals force of attraction
- (4) formation of intermolecular H-bonding.

Ans. (4)





Sol. Carboxylic acid has higher boiling point than aldehyde, ketone and even alcohols of comparable molecular mass.

This is due to more extensive association through intermolecular H-bonding.





53. Compound A,  $\text{C}_8\text{H}_{10}\text{O}$ , is found to react with NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell.

A and Y are respectively



Ans. (3)

Sol. Haloform reaction is shown by compound having



54. The correct difference between first- and second-order reaction is that

- (1) the rate of a first-order reaction does not depend on reactant concentration; the rate of a second-order reaction does depend on reactant concentrations.
- (2) the half-life of a first-order reaction does not depend on  $[\text{A}]_0$ ; the half-life of a second-order reaction does depend on  $[\text{A}]_0$
- (3) a first-order reaction can be catalyzed; a second-order reaction cannot be catalyzed.
- (4) the rate of a first-order reaction does depend on reactant concentrations; the rate of a second-order reaction does not depend on reactant concentrations

Ans. (2)

Sol.  $(t_{1/2})^{1^{\text{st}} \text{ order}} = \text{Independent of Concentration}$

$$(t_{1/2})^{2^{\text{nd}} \text{ order}} \propto \frac{1}{[\text{A}]_0}$$

55. Among  $\text{CaH}_2$ ,  $\text{BeH}_2$ ,  $\text{BaH}_2$ , the order of ionic character is


- (1)  $\text{BeH}_2 < \text{CaH}_2 < \text{BaH}_2$
- (2)  $\text{CaH}_2 < \text{BeH}_2 < \text{BaH}_2$
- (3)  $\text{BeH}_2 < \text{BaH}_2 < \text{CaH}_2$
- (4)  $\text{BaH}_2 < \text{BeH}_2 < \text{CaH}_2$

Ans. (1)

Sol.  $\text{BeH}_2 < \text{CaH}_2 < \text{BaH}_2$

Smaller the size of cation, more will be its polarising power. Hence  $\text{BeH}_2$  will be least ionic.

56. Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below:






Then the species undergoing disproportionation is:-

- (1)  $\text{BrO}_3^-$
- (2)  $\text{BrO}_4^-$
- (3)  $\text{Br}_2$
- (4)  $\text{HBrO}$

Ans. (4)

Sol. Calculate  $E_{\text{cell}}^\circ$  corresponding to each compound under going disproportionation reaction. The reaction for which  $E_{\text{cell}}^\circ$  comes out +ve is spontaneous.



$$E_{\text{cell}}^\circ = \text{SRP (cathode)} - \text{SRP (Anode)}$$

$$= 1.595 - 1.5$$

$$= 0.095 \text{ V}$$

$$E_{\text{cell}}^\circ > 0 \Rightarrow \Delta G^\circ < 0 \text{ [spontaneous]}$$

57. In which case is the number of molecules of water maximum?

- (1) 18 mL of water
- (2) 0.18 g of water
- (3) 0.00224 L of water vapours at 1 atm and 273 K
- (4)  $10^{-3}$  mol of water

Ans. (1)

Sol. (1) 18 mL water

$$\text{As } d_{\text{H}_2\text{O}} = 1 \text{ g/mL} \quad \text{So } W_{\text{H}_2\text{O}} = 18 \text{ g}$$

$$n_{\text{H}_2\text{O}} = \frac{18}{18} = 1$$

$$\text{molecules} = 1 \times N_A$$

(2) 0.18 g of water

$$n_{\text{H}_2\text{O}} = \frac{0.18}{18} = 0.01$$

$$(\text{molecules})_{\text{H}_2\text{O}} = 0.01 \times N_A$$

(3)  $(V_{\text{H}_2\text{O(g)}})_{\text{STP}} = 0.00224 \text{ L}$

$$n_{\text{H}_2\text{O}} = \frac{V}{22.4} = \frac{0.00224}{22.4} = 0.0001$$

$$\text{molecules} = 0.0001 \times N_A$$

(4)  $n_{\text{H}_2\text{O}} = 10^{-3}$

$$(\text{molecules})_{\text{H}_2\text{O}} = 10^{-3} \times N_A$$

58. Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is  $1s^2 2s^2 2p^3$ , the simplest formula for this compound is


- (1)  $\text{Mg}_2\text{X}_3$
- (2)  $\text{MgX}_2$
- (3)  $\text{Mg}_2\text{X}$
- (4)  $\text{Mg}_3\text{X}_2$

Ans. (4)

Sol. Magnesium ion =  $\text{Mg}^{+2}$

X = Nitrogen

Nitrogen ion =  $\text{N}^{-3}$



59. Iron exhibits bcc structure at room temperature. Above  $900^\circ\text{C}$ , it transforms to fcc structure. The ratio of density of iron at room temperature to that at  $900^\circ\text{C}$  (assuming molar mass and atomic radii of iron remains constant with temperature) is

- (1)  $\frac{\sqrt{3}}{\sqrt{2}}$
- (2)  $\frac{4\sqrt{3}}{3\sqrt{2}}$
- (3)  $\frac{3\sqrt{3}}{4\sqrt{2}}$
- (4)  $\frac{1}{2}$

Ans. (3)

Sol. BCC

FCC

$$4r = \sqrt{3}a$$

$$4r = \sqrt{2}a$$


$$a = \frac{4r}{\sqrt{3}}$$

$$a = \frac{4r}{\sqrt{2}}$$

$$\frac{d_{\text{BCC}}}{d_{\text{FCC}}} = \frac{\frac{Z_{\text{BCC}} \times M}{N_A a^3}}{\frac{Z_{\text{FCC}} \times M}{N_A a^3}} = \frac{2 \times M}{N_A \left(\frac{4r}{\sqrt{3}}\right)^3} = \frac{3\sqrt{3}}{4\sqrt{2}}$$

60. Which one is a **wrong** statement ?

- (1) Total orbital angular momentum of electron in 's' orbital is equal to zero
- (2) An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers.
- (3) The electronic configuration of N atom is



(4) The value of m for  $d_{z^2}$  is zero

Ans. (3)

**Sol.** The correct configuration of 'N' is



**61.** Consider the following species:



Which one of these will have the highest bond order?

- (1) NO (2) CN<sup>-</sup>  
(3) CN<sup>+</sup> (4) CN

**Ans. (2)**

Ion/Species	Total electron	Bond order
NO	15	2.5
CN <sup>-</sup>	14	3
CN <sup>+</sup>	12	2
CN	13	2.5

**62.** Which of the following statements is **not** true for halogens ?


- (1) All form monobasic oxyacids.  
(2) All are oxidizing agents.  
(3) All but fluorine show positive oxidation states.  
(4) Chlorine has the highest electron-gain enthalpy.

**Ans. (Bonus)**

**63.** Which one of the following elements is unable to form MF<sub>6</sub><sup>3-</sup> ion ?

- (1) Ga (2) Al (3) B (4) In

**Ans. (3)**




Boron belongs to 2<sup>nd</sup> period and it does not have vacant d-orbital.

**64.** In the structure of ClF<sub>3</sub>, the number of lone pairs of electrons on central atom 'Cl' is

- (1) one (2) two (3) four (4) three

**Ans. (2)**



2 lone pair at equatorial position.

**65.** Considering Ellingham diagram, which of the following metals can be used to reduce alumina ?

- (1) Fe (2) Zn (3) Mg (4) Cu

**Ans. (3)**

**Sol.** Mg has more -ΔG value than alumina. So it will be in the lower part of Ellingham diagram. Metals which has more -ΔG value can reduce those metals oxide which has less -ΔG value.

**66.** The correct order of atomic radii in group 13 elements is

- (1) B < Al < In < Ga < Tl  
(2) B < Al < Ga < In < Tl  
(3) B < Ga < Al < Tl < In  
(4) B < Ga < Al < In < Tl


**Ans. (4)**

**Sol.** In group 13 due to transition contraction [Al > Ga]

**67.** The correct order of N-compounds in its decreasing order of oxidation states is

- (1) HNO<sub>3</sub>, NO, N<sub>2</sub>, NH<sub>4</sub>Cl  
(2) HNO<sub>3</sub>, NO, NH<sub>4</sub>Cl, N<sub>2</sub>  
(3) HNO<sub>3</sub>, NH<sub>4</sub>Cl, NO, N<sub>2</sub>  
(4) NH<sub>4</sub>Cl, N<sub>2</sub>, NO, HNO<sub>3</sub>

**Ans. (1)**



**68.** On which of the following properties does coagulating power of an ion depend ?

- (1) The magnitude of the charge on the alone  
(2) Size of the ion alone  
(3) Both magnitude and sign of the charge the ion  
(4) The sign of charge on the ion alone

**Ans. (3)**

**Sol.** According to Hardy Schulze rule : The coagulating power of an ion depend on both magnitude and sign of the charge of the ion.

**69.** Following solutions were prepared by mixing different volumes of NaOH and HCl of different concentrations :

a. 60mL  $\frac{M}{10}$  HCl + 40mL  $\frac{M}{10}$  NaOH

b. 55mL  $\frac{M}{10}$  HCl + 45mL  $\frac{M}{10}$  NaOH

c. 75mL  $\frac{M}{5}$  HCl + 25mL  $\frac{M}{5}$  NaOH

d. 100mL  $\frac{M}{10}$  HCl + 100mL  $\frac{M}{10}$  NaOH

pH of which one of them will be equal to 1 ?

- (1) b (2) a  
(3) d (4) c

**Ans. (4)**

**Sol.** As N<sub>1</sub>V<sub>1</sub> > N<sub>2</sub>V<sub>2</sub>

So acid is left at the end of reaction

$$N_{\text{final solution}} = [\text{H}^+] = \frac{N_1 V_1 - N_2 V_2}{V_1 + V_2}$$

$$= \frac{\frac{1}{5} \times 75 - \frac{1}{5} \times 25}{75 + 25}$$

$$= \frac{1}{10} = 0.1$$

$$\text{pH} = -\log[\text{H}^+] = 1$$

**70.** The solubility of  $\text{BaSO}_4$  in water  $2.42 \times 10^3 \text{ gL}^{-1}$  at 298 K. The value of solubility product ( $K_{\text{sp}}$ ) will be

(Given molar mass of  $\text{BaSO}_4 = 233 \text{ g mol}^{-1}$ )

- (1)  $1.08 \times 10^{-10} \text{ mol}^2 \text{ L}^{-2}$
- (2)  $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$
- (3)  $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$
- (4)  $1.08 \times 10^{-8} \text{ mol}^2 \text{ L}^{-2}$

**Ans. (1)**

**Sol.** solubility of  $\text{BaSO}_4 = 2.42 \times 10^{-3} \text{ gL}^{-1}$

$$\therefore s = \frac{2.42 \times 10^{-3}}{233} = 1.038 \times 10^{-5} \text{ mol L}^{-1}$$

$$K_{\text{sp}} = s^2 = (1.038 \times 10^{-5})^2 = 1.08 \times 10^{-10} \text{ mol}^2 \text{ L}^{-2}$$

**71.** Given van der Waals constant for  $\text{NH}_3$ ,  $\text{H}_2$  and  $\text{CO}_2$  are respectively 4.17, 0.244, 1.36 and 3.59, which one of the following gases is most easily liquefied?

- (1)  $\text{NH}_3$
- (2)  $\text{H}_2$
- (3)  $\text{O}_2$
- (4)  $\text{CO}_2$




**Ans. (1)**

**Sol.** Critical temperature  $\propto$  vanderwaal constant (a) maximum "a"  $\Rightarrow$  gas with maximum  $T_C \Rightarrow$  easiest liquification =  $\text{NH}_3$

**72.** The compound A on treatment with Na gives B, and with  $\text{PCl}_5$  gives C. B and C react together to give diethyl ether. A, B and C are in the order

- (1)  $\text{C}_2\text{H}_5\text{OH}$ ,  $\text{C}_2\text{H}_6$ ,  $\text{C}_2\text{H}_5\text{Cl}$
- (2)  $\text{C}_2\text{H}_5\text{OH}$ ,  $\text{C}_2\text{H}_5\text{Cl}$ ,  $\text{C}_2\text{H}_5\text{ONa}$
- (3)  $\text{C}_2\text{H}_5\text{Cl}$ ,  $\text{C}_2\text{H}_6$ ,  $\text{C}_2\text{H}_5\text{OH}$
- (4)  $\text{C}_2\text{H}_5\text{OH}$ ,  $\text{C}_2\text{H}_5\text{ONa}$ ,  $\text{C}_2\text{H}_5\text{Cl}$


**Ans. (4)**



**73.** Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms. (A) is

- (1)  $\text{CH}\equiv\text{CH}$
- (2)  $\text{CH}_2=\text{CH}_2$
- (3)  $\text{CH}_3-\text{CH}_3$
- (4)  $\text{CH}_4$

**Ans. (4)**



**74.** The compound  $\text{C}_7\text{H}_8$  undergoes the following reactions :




The product 'C' is

- (1) m-bromotoluene
- (2) o-bromotoluene
- (3) 3-bromo-2,4,6-trichlorotoluene
- (4) p-bromotoluene

**Ans. (1)**

**Sol.**



**75.** Which oxide of nitrogen is **not** a common pollutant introduced into the atmosphere both due to natural and human activity ?

- (1)  $\text{N}_2\text{O}_5$
- (2)  $\text{NO}_2$
- (3)  $\text{N}_2\text{O}$
- (4)  $\text{NO}$

**Ans. (1)**

**Sol.** Nitrous oxide ( $\text{N}_2\text{O}$ ) occurs naturally in environment. In automobile engine, when fossil is burnt dinitrogen & dioxygen combine to yield  $\text{NO}$  &  $\text{NO}_2$ .



**76.** For the redox reaction



the correct coefficients of the reactants for the balanced equation are

- |     | $\text{MnO}_4^-$ | $\text{C}_2\text{O}_4^{2-}$ | $\text{H}^+$ |
|-----|------------------|-----------------------------|--------------|
| (1) | 16               | 5                           | 2            |
| (2) | 2                | 5                           | 16           |
| (3) | 2                | 16                          | 5            |
| (4) | 5                | 16                          | 2            |

**Ans. (2)**




multiplying (1) by 2 and (2) by 5 to balance  $e^-$



on balancing charge;



77. Which one of the following conditions will favour maximum formation of the product in the reaction,



- (1) Low temperature and high pressure
- (2) Low temperature and low pressure
- (3) High temperature and high pressure
- (4) High temperature and low pressure

Ans. (1)

Sol. For reaction  $\Delta H = -ve$  and  $\Delta n_g = -ve$   
 $\therefore$  High P, Low T, favour product formation.

78. The correction factor 'a' to the ideal gas equation corresponds to

- (1) density of the gas molecules
- (2) volume of the gas molecules
- (3) electric field present between the gas molecules
- (4) forces of attraction between the gas molecules

Ans. (4)

Sol. Vanderwaal constant (a)  $\propto$  forces of attraction.

79. When initial concentration of the reactant is doubled, the half-life period of a zero order reaction

- (1) is halved
- (2) is doubled
- (3) is tripled
- (4) remains unchanged

Ans. (2)

Sol.  $(t_{1/2})_{\text{zero}} = \frac{[A]_0}{2K}$


$\therefore$  If  $[A]_0 = \text{doubled}$ ,  $t_{1/2} = \text{doubled}$

80. The bond dissociation energies of  $X_2$ ,  $Y_2$  and  $XY$  are in the ratio of 1 : 0.5 : 1.  $\Delta H$  for the formation of  $XY$  is  $-200 \text{ kJ mol}^{-1}$ . The bond dissociation energy of  $X_2$  will be

- (1)  $200 \text{ kJ mol}^{-1}$
- (2)  $100 \text{ kJ mol}^{-1}$
- (3)  $800 \text{ kJ mol}^{-1}$
- (4)  $400 \text{ kJ mol}^{-1}$

Ans. (3)

Sol. let B.E. of  $x_2$ ,  $y_2$  &  $xy$  are  $x \text{ kJ mol}^{-1}$ ,  $0.5x \text{ kJ mol}^{-1}$  and  $x \text{ kJ mol}^{-1}$  respectively




$$\Delta H = -200 = \Sigma(\text{B.E})_{\text{Reactant}} - \Sigma(\text{B.E})_{\text{Product}}$$

$$= \left[ \frac{1}{2} \times (x) + \frac{1}{2} \times (0.5x) \right] - [1 \times (x)]$$

$$\text{B.E of } X_2 = x = 800 \text{ kJ mol}^{-1}$$




81. Identify the major products P, Q and R in the following sequence of reaction :



- |     |   |   |   |   |                                    |
|-----|---|---|---|---|------------------------------------|
|     | P |   | Q |   | R                                  |
| (1) |   | , |   | , | $\text{CH}_3\text{CH}_2\text{-OH}$ |
| (2) |   | , |   | , |                                    |
| (3) |   | , |   | , | $\text{CH}_3\text{CH(OH)CH}_3$     |
| (4) |   | , |   | , | $\text{CH}_3\text{-CO-CH}_3$       |

Ans. (4)

Sol.




**82.** Which of the following compounds can form a zwitterion ?

- (1) Aniline (2) Acetanilide  
(3) Benzoic acid (4) Glycine

**Ans. (4)**


**Sol.** The molecule which forms zwitter ion is glycine.



**83.** The type of isomerism shown by the complex  $[\text{CoCl}_2(\text{en})_2]$  is

- (1) Geometrical isomerism  
(2) Coordination isomerism  
(3) Ionization isomerism  
(4) Linkage isomerism

**Ans. (1)**



**84.** Which one of the following ions exhibits d-d transition and paramagnetism as well ?

- (1)  $\text{CrO}_4^{2-}$  (2)  $\text{Cr}_2\text{O}_7^{2-}$   
(3)  $\text{MnO}_4^-$  (4)  $\text{MnO}_4^{2-}$

**Ans. (4)**


**Sol.**  $\text{CrO}_4^{2-}$   $\text{Cr}^{+6}$  diamagnetic

$\text{Cr}_2\text{O}_7^{2-}$   $\text{Cr}^{+6}$  diamagnetic

$\text{MnO}_4^-$   $\text{Mn}^{+7}$  diamagnetic

$\text{MnO}_4^{2-}$   $\text{Mn}^{+6}$  paramagnetic 1          

unpaired electron is present so d-d transition is possible.





**85.** The geometry and magnetic behaviour of the complex  $[\text{Ni}(\text{CO})_4]$  are


- (1) square planar geometry and diamagnetic  
(2) tetrahedral geometry and diamagnetic  
(3) square planar geometry and paramagnetic  
(4) tetrahedral geometry and paramagnetic

**Ans. (2)**

**Sol.** tetrahedral geometry and diamagnetic



CO is SFL so unpaired electrons will get paired.



$sp^3$  hybridisation

Tetrahedral, diamagnetic

**86.** Iron carbonyl,  $\text{Fe}(\text{CO})_5$  is

- (1) tetranuclear (2) mononuclear  
(3) trinuclear (4) dinuclear

**Ans. (2)**

**Sol.**  $\text{Fe}(\text{CO})_5$

EAN = Z-O.N. + 2(C.N.)

$$= 26 - 0 + 2(5)$$

$$= 26 + 10$$

$$= 36$$

only one central metal atom/ion is present and it follows EAN rule, so it is mononuclear



**87.** Match the metal ions given in Column I with the spin magnetic moments of the ions given in Column II and assign the **correct** code :

Column I	Column II
a. $\text{Co}^{3+}$	i. $\sqrt{8}$ B.M.
b. $\text{Cr}^{3+}$	ii. $\sqrt{35}$ B.M.
c. $\text{Fe}^{3+}$	iii. $\sqrt{3}$ B.M.
d. $\text{Ni}^{2+}$	iv. $\sqrt{24}$ B.M.
	v. $\sqrt{15}$ B.M.

	a	b	c	d
(1)	iv	v	ii	i
(2)	i	ii	iii	iv
(3)	iv	i	ii	iii
(4)	iii	v	i	ii


**Ans. (1)**

**Sol.** Magnetic moment ( $\mu$ ) =  $\sqrt{n(n+2)}$  B.M.





$n = 4$

$\mu = \sqrt{4(4+2)} = \sqrt{24}$  B. M





$n = 3$

$\mu = \sqrt{3(3+2)} = \sqrt{15}$  B.M.



$n = 5$

$\mu = \sqrt{5(5+2)} = \sqrt{35}$  B. M.



$n = 2$

$\mu = \sqrt{2(2+2)} = \sqrt{8}$  B. M.

**88.** Which of the following is correct with respect to -I effect of the substituents ? (R = alkyl)



- (1)  $-\text{NH}_2 < -\text{OR} < -\text{F}$
- (2)  $-\text{NR}_2 < -\text{OR} < -\text{F}$
- (3)  $-\text{NH}_2 > -\text{OR} > -\text{F}$
- (4)  $-\text{NR}_2 > -\text{OR} > -\text{F}$

**Ans. (1/2)**

**Sol.** (Based on EN)




**89.** Which of the following carbocations is expected to be most stable ?




**Ans. (3)**

**Sol.**  $-\text{NO}_2$  group is meta-directing group



(Less stable due to more  $e^-$  withdrawing effect of  $-\text{NO}_2$ )



(More stable due to less  $e^-$  withdrawing effect of  $-\text{NO}_2$ )

**90.** Which of the following molecules represents the order of hybridisation  $sp^2$ ,  $sp^2$ ,  $sp$ ,  $sp$  from left to right atoms ?

- (1)  $\text{HC} \equiv \text{C} - \text{C} \equiv \text{CH}$
- (2)  $\text{CH}_2 = \text{CH} - \text{C} \equiv \text{CH}$
- (3)  $\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$
- (4)  $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_3$

**Ans. (2)**

